Food and Drug Administration, HHS

(3) Limitations. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[69 FR 47363, Aug. 5, 2004, as amended at 77 FR 64717, Oct. 23, 2012]

§ 524.1044d Gentamicin and betamethasone ointment.

- (a) Specifications. Each gram of ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin base and betamethasone valerate equivalent to 1 milligram of betamethasone.
- (b) *Sponsor*. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use in dogs—(1) Amount—(i) Otitis externa. Instill 3 to 8 drops into the ear canal twice daily for 7 days.
- (ii) Infected superficial lesions. Apply to cover the treatment area twice daily for 7 to 14 days.
- (2) Indications for use. For the treatment of acute and chronic otitis externa and infected superficial lesions caused by bacteria sensitive to gentamicin.
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 26378, June 18, 1982, as amended at 52 FR 7832, Mar. 13, 1987; 79 FR 10969, Feb. 27, 2014]

§524.1044e Gentamicin spray.

- (a) Specification. Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to 1.07 milligrams of gentamicin.
- (b) Sponsor. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use in cattle—(1) Amount. Hold the sprayer upright 3 to 6 inches from the affected eye, with the opening directed towards the eye, and pump once. Treat once daily for up to 3 days.
- (2) *Indications for use*. For the treatment of pinkeye in cattle (infectious bovine keratoconjunctivitis) caused by *Moraxella bovis*.
- (3) Limitations. Conditions other than bacterial infections of the bovine eye and infectious keratoconjunctivitis caused by Moraxella bovis may produce similar signs. If conditions persists or

increases, discontinue use and consult a veterinarian.

 $[48\ FR\ 41157,\ Sept.\ 14,\ 1983,\ as\ amended\ 52\ FR\ 7833,\ Mar.\ 13,\ 1987;\ 79\ FR\ 10969,\ Feb.\ 27,\ 2014]$

§ 524.1044f Gentamicin and betamethasone spray.

- (a) Specifications. Each milliliter of spray contains gentamicin sulfate equivalent to 0.57 milligram (mg) gentamicin base and betamethasone valerate equivalent to 0.284 mg betamethasone.
- (b) Sponsors. See Nos. 000061, 054925, 058005, 058829, and 065531 in $\S510.600(c)$ of this chapter.
- (c) Conditions of use in dogs—(1) Amount. Hold bottle upright 3 to 6 inches from the lesion and depress the sprayer head twice. Administer two spray actuations two to four times daily for 7 days.
- (2) Indications for use. For the treatment of infected superficial lesions caused by bacteria susceptible to gentamicin.
- (3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[71 FR 13542, Mar. 16, 2006, as amended at 72 FR 5929, Feb. 8, 2007; 74 FR 22822, May 15, 2009; 77 FR 3598, Jan. 25, 2012]

§ 524.1044g Gentamicin, betamethasone, and clotrimazole ointment.

- (a) Specifications. Each gram (g) of ointment contains gentamicin sulfate equivalent to 3 milligrams (mg) gentamicin base, betamethasone valerate equivalent to 1 mg betamethasone, and 10 mg clotrimazole.
- (b) *Sponsors*. See sponsors in §510.600(c) of this chapter for uses as in paragraph (c) of this section.
- (1) No. 000061 for use of 7.5- or 15-gram (g) tubes, 12.5-, 30-, or 215-g bottles.
- (2) No. 054925 for use of 7.5- or 15-g tubes; 10-, 15-, 25-, or 215-g bottles.
- (3) No. 000859 for use of 10-, 20-, 40-, or 215-g bottles.
- (4) No. 025463 for use of 7.5- or 15-g tubes, or 215-g bottles.
- (c) Conditions of use in dogs—(1) Amount. Instill ointment twice daily into the ear canal for 7 consecutive days.